

# Undecidability of the word problem for one-relator inverse monoids

Robert D. Gray<sup>1</sup>

SandGAL 2019, Cremona, Italy, June 2019



---

<sup>1</sup>Research supported by the EPSRC grant EP/N033353/1 "Special inverse monoids: subgroups, structure, geometry, rewriting systems and the word problem".

## Word problem for one-relator groups and monoids

	$\text{Gp}\langle A \mid w = 1 \rangle$ $\text{FG}(A)/\langle\langle w \rangle\rangle$	$\text{Mon}\langle A \mid w = 1 \rangle$ $A^*/\langle\langle (w, 1) \rangle\rangle$	$\text{Inv}\langle A \mid w = 1 \rangle$ $\text{FIM}(A)/\langle\langle (w, 1) \rangle\rangle$
Word problem decidable	Magnus (1932) ✓	Adjan (1966) ✓	?

Theorem (Scheiblich (1973) & Munn (1974))

Free inverse monoids have decidable word problem.

Conjecture (Margolis, Meakin, Stephen (1987))

If  $M = \text{Inv}\langle A \mid w = 1 \rangle$ , then the word problem for  $M$  is decidable.

## Word problem for one-relator groups and monoids

	$\text{Gp}\langle A \mid w = 1 \rangle$ $\text{FG}(A)/\langle\langle w \rangle\rangle$	$\text{Mon}\langle A \mid w = 1 \rangle$ $A^*/\langle\langle (w, 1) \rangle\rangle$	$\text{Inv}\langle A \mid w = 1 \rangle$ $\text{FIM}(A)/\langle\langle (w, 1) \rangle\rangle$
Word problem decidable	Magnus (1932) ✓	Adjan (1966) ✓	?

### Theorem (Scheiblich (1973) & Munn (1974))

Free inverse monoids have decidable word problem.

### Conjecture (Margolis, Meakin, Stephen (1987))

If  $M = \text{Inv}\langle A \mid w = 1 \rangle$ , then the word problem for  $M$  is decidable.

### Theorem (Ivanov, Margolis, Meakin (2001))

If the word problem is decidable for all inverse monoids of the form  $\text{Inv}\langle A \mid w = 1 \rangle$  then the word problem is also decidable for every one-relator monoid  $\text{Mon}\langle A \mid u = v \rangle$ .

## Word problem for one-relator groups and monoids

	$\text{Gp}\langle A \mid w = 1 \rangle$ $\text{FG}(A)/\langle\langle w \rangle\rangle$	$\text{Mon}\langle A \mid w = 1 \rangle$ $A^*/\langle\langle (w, 1) \rangle\rangle$	$\text{Inv}\langle A \mid w = 1 \rangle$ $\text{FIM}(A)/\langle\langle (w, 1) \rangle\rangle$
Word problem decidable	Magnus (1932) ✓	Adjan (1966) ✓	?

### Theorem (Scheiblich (1973) & Munn (1974))

Free inverse monoids have decidable word problem.

### Conjecture (Margolis, Meakin, Stephen (1987))

If  $M = \text{Inv}\langle A \mid w = 1 \rangle$ , then the word problem for  $M$  is decidable.

Proved true in many cases e.g. when  $w$  satisfies...

- ▶ Idempotent word [Birget, Margolis, Meakin, 1993, 1994]
- ▶  $w$ -strictly positive [Ivanov, Margolis, Meakin, 2001]
- ▶ Adjan or Baumslag-Solitar type [Margolis, Meakin, Šuník, 2005]
- ▶ Sparse word [Hermiller, Lindblad, Meakin, 2010]
- ▶ Certain small cancellation conditions [A. Juhász, 2012, 2014]

## Word problem for one-relator groups and monoids

	$\text{Gp}\langle A \mid w = 1 \rangle$ $\text{FG}(A)/\langle\langle w \rangle\rangle$	$\text{Mon}\langle A \mid w = 1 \rangle$ $A^*/\langle\langle (w, 1) \rangle\rangle$	$\text{Inv}\langle A \mid w = 1 \rangle$ $\text{FIM}(A)/\langle\langle (w, 1) \rangle\rangle$
Word problem decidable	Magnus (1932) ✓	Adjan (1966) ✓	✗

### Theorem (RDG (2019))

There is a one-relator inverse monoid  $\text{Inv}\langle A \mid w = 1 \rangle$  with undecidable word problem.

## Word problem for one-relator groups and monoids

	$\text{Gp}\langle A \mid w = 1 \rangle$ $\text{FG}(A)/\langle\langle w \rangle\rangle$	$\text{Mon}\langle A \mid w = 1 \rangle$ $A^*/\langle\langle (w, 1) \rangle\rangle$	$\text{Inv}\langle A \mid w = 1 \rangle$ $\text{FIM}(A)/\langle\langle (w, 1) \rangle\rangle$
Word problem decidable	Magnus (1932) ✓	Adjan (1966) ✓	✗

### Theorem (RDG (2019))

There is a one-relator inverse monoid  $\text{Inv}\langle A \mid w = 1 \rangle$  with undecidable word problem.

#### Ingredients for the proof:

- ▶ Submonoid membership problem for one relator groups.
- ▶ HNN-extensions and free products of groups.
- ▶ Right-angled Artin groups (RAAGs).
- ▶ Right units of special inverse monoids

$$\text{Inv}\langle A \mid w_1 = 1, w_2 = 1, \dots, w_k = 1 \rangle$$

and Stephen's procedure for constructing Schützenberger graphs.

- ▶ Properties of  $E$ -unitary inverse monoids.

# Inverse monoid presentations

An **inverse monoid** is a monoid  $M$  such that for every  $x \in M$  there is a unique  $x^{-1} \in M$  such that  $xx^{-1}x = x$  and  $x^{-1}xx^{-1} = x^{-1}$ .


For all  $x, y \in M$  we have

$$x = xx^{-1}x, (x^{-1})^{-1} = x, (xy)^{-1} = y^{-1}x^{-1}, xx^{-1}yy^{-1} = yy^{-1}xx^{-1} \quad (\dagger)$$

$$\text{Inv}\langle A \mid u_i = v_i \ (i \in I) \rangle = \text{Mon}\langle A \cup A^{-1} \mid u_i = v_i \ (i \in I) \cup (\dagger) \rangle$$

where  $u_i, v_i \in (A \cup A^{-1})^*$  and  $x, y$  range over all words from  $(A \cup A^{-1})^*$ .

**Free inverse monoid**  $\text{FIM}(A) = \text{Inv}\langle A \mid \rangle$



**Munn (1974)**

Elements of  $\text{FIM}(A)$  can be represented using Munn trees. e.g. in  $\text{FIM}(a, b)$  we have  $u = w$  where

$$u = aa^{-1}bb^{-1}ba^{-1}abb^{-1}$$

$$w = bbb^{-1}a^{-1}ab^{-1}aa^{-1}b$$

# The word problem

$M$  - a finitely generated monoid with a finite generating set  $A$ .

$\pi : A^* \rightarrow M$  - the canonical monoid homomorphism.

The monoid  $M$  has decidable word problem if there is an algorithm which solves the following decision problem:

**INPUT:** Two words  $u, v \in A^*$ .

**QUESTION:**  $\pi(u) = \pi(v)$ ? i.e. do  $u$  and  $v$  represent the same element of the monoid  $M$ ?


For a group or an inverse monoid with generating set  $A$  the word problem is defined in the same way except the input is two words  $u, v \in (A \cup A^{-1})^*$ .

**Example.** The bicyclic monoid  $\text{Inv}\langle a \mid aa^{-1} = 1 \rangle$  has decidable word problem.



## Proof strategy

$$M = \text{Inv}\langle A \mid r=1 \rangle \longrightarrow G = \text{Gp}\langle A \mid r=1 \rangle$$



If  $M$  has decidable word problem

$\Rightarrow$  membership problem for  $U_R \leq M$  is decidable

since for  $w \in (A \cup A^{-1})^*$

$$w \in U_R \iff ww^{-1} = 1$$

(sometimes)

$\rightsquigarrow$  membership problem for  $N \leq G$  is decidable

# RAAGs induced subgraphs and subgroups


## Definition

The **right-angled Artin group**  $A(\Gamma)$  associated with the graph  $\Gamma$  is

$$\text{Gp}\langle V\Gamma \mid uv = vu \text{ if and only if } \{u, v\} \in E\Gamma \rangle.$$

**Fact:** If  $\Delta$  is an induced subgraph of  $\Gamma$  then the embedding  $\Delta \rightarrow \Gamma$  induces an embedding  $A(\Delta) \rightarrow A(\Gamma)$ .

## Example




$$\begin{array}{l} A(\Delta) = \text{Gp} \langle a, c, d, e \mid ac = ca, de = ed \rangle \\ \hookrightarrow A(\Gamma) = \text{Gp} \langle a, b, c, d, e \mid ac = ca, de = ed, \\ ab = ba, bc = cb, \\ bd = db \rangle \end{array}$$

## HNN-extensions of groups


$H \cong \text{Gp}\langle A \mid R \rangle$ ,  $K, L \leq H$  with  $K \cong L$ . Let  $\phi : K \rightarrow L$  be an isomorphism. The **HNN-extension** of  $H$  with respect to  $\phi$  is

$$G = \text{HNN}(H, \phi) = \text{Gp}\langle A, t \mid R, t^{-1}kt = \phi(k) \ (k \in K) \rangle$$

**Fact:**  $H$  embeds naturally into the HNN extension  $G = \text{HNN}(H, \phi)$ .



# HNN-extensions of RAAGs



## Definition

$\Gamma$  - finite graph,  $\psi : \Delta_1 \rightarrow \Delta_2$  an isomorphism between finite induced subgraphs.

$A(\Gamma, \psi)$  is defined to be the HNN-extension of  $A(\Gamma)$  with respect to the isomorphism  $A(\Delta_1) \rightarrow A(\Delta_2)$  induced by  $\psi$ .

**Fact:**  $A(\Gamma)$  embeds naturally into  $A(\Gamma, \psi)$ .

## HNN-extension of $A(P_4)$ over $A(P_3)$

Let  $P_4$  be the graph




$$A(P_4) = \text{Gp}\langle a, b, c, d \mid ab = ba, bc = cb, cd = dc \rangle.$$

$\Delta_1$  - subgraph induced by  $\{a, b, c\}$ ,  $\Delta_2$  subgraph induced by  $\{b, c, d\}$ ,  
 $\psi : \Delta_1 \rightarrow \Delta_2$  - the isomorphism  $a \mapsto b$ ,  $b \mapsto c$ , and  $c \mapsto d$ .

## HNN-extension of $A(P_4)$ over $A(P_3)$

Let  $P_4$  be the graph



$$A(P_4) = \text{Gp}\langle a, b, c, d \mid ab = ba, bc = cb, cd = dc \rangle.$$

$\Delta_1$  - subgraph induced by  $\{a, b, c\}$ ,  $\Delta_2$  subgraph induced by  $\{b, c, d\}$ ,


$\psi : \Delta_1 \rightarrow \Delta_2$  - the isomorphism  $a \mapsto b, b \mapsto c$ , and  $c \mapsto d$ .

Then the HNN-extension  $A(P_4, \psi)$  of  $A(P_4)$  with respect to  $\psi$  is

$$\begin{aligned} & A(P_4, \psi) \\ = & \text{Gp}\langle a, b, c, d, t \mid ab = ba, bc = cb, cd = dc, tat^{-1} = b, tbt^{-1} = c, tct^{-1} = d \rangle \end{aligned}$$

## HNN-extension of $A(P_4)$ over $A(P_3)$

Let  $P_4$  be the graph



$$A(P_4) = \text{Gp}\langle a, b, c, d \mid ab = ba, bc = cb, cd = dc \rangle.$$

$\Delta_1$  - subgraph induced by  $\{a, b, c\}$ ,  $\Delta_2$  subgraph induced by  $\{b, c, d\}$ ,


$\psi : \Delta_1 \rightarrow \Delta_2$  - the isomorphism  $a \mapsto b$ ,  $b \mapsto c$ , and  $c \mapsto d$ .

Then the HNN-extension  $A(P_4, \psi)$  of  $A(P_4)$  with respect to  $\psi$  is

$$\begin{aligned} & A(P_4, \psi) \\ = & \text{Gp}\langle a, b, c, d, t \mid ab = ba, bc = cb, cd = dc, tat^{-1} = b, tbt^{-1} = c, tct^{-1} = d \rangle \\ = & \text{Gp}\langle a, t \mid a(tat^{-1}) = (tat^{-1})a, (tat^{-1})(t^2at^{-2}) = (t^2at^{-2})(tat^{-1}), \\ & (t^2at^{-2})(t^3at^{-3}) = (t^3at^{-3})(t^2at^{-2}) \rangle. \end{aligned}$$

## HNN-extension of $A(P_4)$ over $A(P_3)$

Let  $P_4$  be the graph



$$A(P_4) = \text{Gp}\langle a, b, c, d \mid ab = ba, bc = cb, cd = dc \rangle.$$

$\Delta_1$  - subgraph induced by  $\{a, b, c\}$ ,  $\Delta_2$  subgraph induced by  $\{b, c, d\}$ ,

$\psi : \Delta_1 \rightarrow \Delta_2$  - the isomorphism  $a \mapsto b$ ,  $b \mapsto c$ , and  $c \mapsto d$ .


Then the HNN-extension  $A(P_4, \psi)$  of  $A(P_4)$  with respect to  $\psi$  is

$$\begin{aligned} & A(P_4, \psi) \\ = & \text{Gp}\langle a, b, c, d, t \mid ab = ba, bc = cb, cd = dc, tat^{-1} = b, tbt^{-1} = c, tct^{-1} = d \rangle \\ = & \text{Gp}\langle a, t \mid a(tat^{-1}) = (tat^{-1})a, (tat^{-1})(t^2at^{-2}) = (t^2at^{-2})(tat^{-1}), \\ & (t^2at^{-2})(t^3at^{-3}) = (t^3at^{-3})(t^2at^{-2}) \rangle. \\ = & \text{Gp}\langle a, t \mid atat^{-1}a^{-1}ta^{-1}t^{-1} = 1 \rangle. \end{aligned}$$



## HNN-extension of $A(P_4)$ over $A(P_3)$

Let  $P_4$  be the graph



$$A(P_4) = \text{Gp}\langle a, b, c, d \mid ab = ba, bc = cb, cd = dc \rangle.$$

$\Delta_1$  - subgraph induced by  $\{a, b, c\}$ ,  $\Delta_2$  subgraph induced by  $\{b, c, d\}$ ,

$\psi : \Delta_1 \rightarrow \Delta_2$  - the isomorphism  $a \mapsto b$ ,  $b \mapsto c$ , and  $c \mapsto d$ .

Then the HNN-extension  $A(P_4, \psi)$  of  $A(P_4)$  with respect to  $\psi$  is

$$\begin{aligned} & A(P_4, \psi) \\ = & \text{Gp}\langle a, b, c, d, t \mid ab = ba, bc = cb, cd = dc, tat^{-1} = b, tbt^{-1} = c, tct^{-1} = d \rangle \\ = & \text{Gp}\langle a, t \mid a(tat^{-1}) = (tat^{-1})a, (tat^{-1})(t^2at^{-2}) = (t^2at^{-2})(tat^{-1}), \\ & (t^2at^{-2})(t^3at^{-3}) = (t^3at^{-3})(t^2at^{-2}) \rangle. \\ = & \text{Gp}\langle a, t \mid atat^{-1}a^{-1}ta^{-1}t^{-1} = 1 \rangle. \end{aligned}$$

### Conclusion

$A(P_4)$  embeds into the one-relator group

$$A(P_4, \psi) = \text{Gp}\langle a, t \mid atat^{-1}a^{-1}ta^{-1}t^{-1} = 1 \rangle.$$

# Submonoid membership problem

$G$  - a finitely generated group with a finite group generating set  $A$ .

$\pi : (A \cup A^{-1})^* \rightarrow G$  - the canonical monoid homomorphism.

$T$  - a finitely generated submonoid of  $G$ .

The **membership problem for  $T$  within  $G$  is decidable** if there is an algorithm which solves the following decision problem:

**INPUT:** A word  $w \in (A \cup A^{-1})^*$ .

**QUESTION:**  $\pi(w) \in T$ ?

## Theorem B

Let  $G$  be the one-relator group  $\text{Gp}\langle a, t \mid atat^{-1}a^{-1}ta^{-1}t^{-1} = 1 \rangle$ . Then there is a fixed finitely generated submonoid  $N$  of  $G$  such that the membership problem for  $N$  within  $G$  is undecidable.


# Proof of Theorem B

## Theorem B


Let  $G$  be the one-relator group  $\text{Gp}\langle a, t \mid atat^{-1}a^{-1}ta^{-1}t^{-1} = 1 \rangle$ . Then there is a fixed finitely generated submonoid  $N$  of  $G$  such that the membership problem for  $N$  within  $G$  is undecidable.

**Proof.** By [Lohrey & Steinberg, 2008] there is a finitely generated submonoid  $T$  of  $A(P_4)$  such that the membership problem for  $T$  within  $A(P_4)$  is undecidable. Let  $\theta : A(P_4) \rightarrow G$  be an embedding. Then  $N = \theta(T)$  is a finitely generated submonoid of  $G$  such that the membership problem for  $N$  within  $G$  is undecidable.

$A(P_4)$




$\text{Gp}\langle a, t \mid atat^{-1}a^{-1}ta^{-1}t^{-1} = 1 \rangle$



## Proof strategy

$$M = \text{Inv}\langle A \mid r=1 \rangle \longrightarrow G = \text{Gp}\langle A \mid r=1 \rangle$$



If  $M$  has decidable word problem

$\Rightarrow$  membership problem for  $U_R \leq M$  is decidable

since for  $w \in (A \cup A^{-1})^*$

$$w \in U_R \iff ww^{-1} = 1$$

(sometimes)

$\rightsquigarrow$  membership problem for  $N \leq G$  is decidable

## General observations about inverse monoids

$S$  – an inverse monoid generated by  $A$ ,  $E(S)$  – set of idempotents,

$U_R \leq S$  – right units = submonoid of right invertible elements.

- ▶ If  $e \in E(S)$  and  $e \in U_R$  then  $e = 1$ .
- ▶ **Two relations for the price of one:** If  $e$  is an idempotent in  $\text{FIM}(A)$  and  $r \in (A \cup A^{-1})^*$  then

$$\text{Inv}\langle A \mid er = 1 \rangle = \text{Inv}\langle A \mid e = 1, r = 1 \rangle.$$

- ▶  $e \in (A \cup A^{-1})^*$  is an idempotent in  $\text{FIM}(A)$  if and only if  $e$  freely reduces to 1 in the free group  $\text{FG}(A)$ . e.g.

$$x^{-1}y^{-1}xx^{-1}yzz^{-1}x \in E(\text{FIM}(x, y, z)).$$

## A general construction

For any  $r, w_1, \dots, w_k \in (A \cup A^{-1})^*$ , with  $A = \{a_1, \dots, a_n\}$ , set  $e$  equal to  $a_1 a_1^{-1} \dots a_n a_n^{-1} (tw_1 t^{-1})(tw_1^{-1} t^{-1})(tw_2 t^{-1})(tw_2^{-1} t^{-1}) \dots (tw_k t^{-1})(tw_k^{-1} t^{-1}) a_n^{-1} a_n \dots a_1^{-1} a_1$

where  $t$  is a new symbol. Then

$$\begin{aligned} M &= \text{Inv}\langle A, t \mid er = 1 \rangle \\ &= \text{Inv}\langle A, t \mid r = 1, aa^{-1} = 1, a^{-1}a = 1 \ (a \in A), (tw_i t^{-1})(tw_i t^{-1})^{-1} = 1 \ (1 \leq i \leq k) \rangle \\ &\cong \text{Gp}\langle A \mid r = 1 \rangle * \text{FIM}(t) / \{(tw_i t^{-1})(tw_i t^{-1})^{-1} = 1 \ (1 \leq i \leq k)\}. \end{aligned}$$

### Key claim

Let  $T$  be the submonoid of  $G = \text{Gp}\langle A \mid r = 1 \rangle$  generated by  $\{w_1, w_2, \dots, w_k\}$ . Then for all  $u \in (A \cup A^{-1})^*$  we have

$$u \in T \text{ in } G \iff tut^{-1} \in U_R \text{ in } M.$$

## A general construction

For any  $r, w_1, \dots, w_k \in (A \cup A^{-1})^*$ , with  $A = \{a_1, \dots, a_n\}$ , set  $e$  equal to  $a_1 a_1^{-1} \dots a_n a_n^{-1} (tw_1 t^{-1})(tw_1^{-1} t^{-1})(tw_2 t^{-1})(tw_2^{-1} t^{-1}) \dots (tw_k t^{-1})(tw_k^{-1} t^{-1}) a_n^{-1} a_n \dots a_1^{-1} a_1$

where  $t$  is a new symbol. Then

$$\begin{aligned} M &= \text{Inv}\langle A, t \mid er = 1 \rangle \\ &= \text{Inv}\langle A, t \mid r = 1, aa^{-1} = 1, a^{-1}a = 1 \ (a \in A), (tw_i t^{-1})(tw_i t^{-1})^{-1} = 1 \ (1 \leq i \leq k) \rangle \\ &\cong \text{Gp}\langle A \mid r = 1 \rangle * \text{FIM}(t) / \{(tw_i t^{-1})(tw_i t^{-1})^{-1} = 1 \ (1 \leq i \leq k)\}. \end{aligned}$$

### Key claim

Let  $T$  be the submonoid of  $G = \text{Gp}\langle A \mid r = 1 \rangle$  generated by  $\{w_1, w_2, \dots, w_k\}$ . Then for all  $u \in (A \cup A^{-1})^*$  we have

$$u \in T \text{ in } G \iff tut^{-1} \in U_R \text{ in } M.$$


### Theorem

If  $M = \text{Inv}\langle A, t \mid er = 1 \rangle$  has decidable word problem then the membership problem for  $T$  within  $G = \text{Gp}\langle A \mid r = 1 \rangle$  is decidable.

## Proof strategy refined

$$M = \text{Inv} \langle A, t \mid e r = 1 \rangle \longrightarrow G_p \langle A, t \mid r = 1 \rangle \\ = G_p \langle A \mid r = 1 \rangle * \text{FG}(t) \\ = G$$

$t u t^{-1} = \gamma$
$U_R = \{ m \in M : m m^{-1} = 1 \}$
$t v t^{-1}$



If  $M$  has decidable word problem

$\Rightarrow$  membership problem for  $U_R \leq M$  is decidable

$\Rightarrow \forall u \in (A \cup A^{-1})^*$  can decide  $t u t^{-1} \in U_R ?$

$\Rightarrow$  (by key claim) can decide  $u \in T = \text{Mon} \langle w_1, \dots, w_k \rangle \leq G$



# Tying things together

## Thorem A

There is a one-relator inverse monoid  $\text{Inv}\langle A \mid w = 1 \rangle$  with undecidable word problem.

## Proof.

Let  $A = \{a, z\}$  and let  $G$  be the one-relator group

$$\text{Gp}\langle a, z \mid aza z^{-1} a^{-1} z a^{-1} z^{-1} = 1 \rangle.$$

Let  $W = \{w_1, \dots, w_k\}$  be a finite subset of  $(A \cup A^{-1})^*$  such that the membership problem for  $T = \text{Mon}\langle W \rangle$  within  $G$  is undecidable. Such a set  $W$  exists by Theorem B. Set  $e$  to be the idempotent word

$$aa^{-1}zz^{-1}(tw_1t^{-1})(tw_1^{-1}t^{-1})(tw_2t^{-1})(tw_2^{-1}t^{-1})\dots(tw_kt^{-1})(tw_k^{-1}t^{-1})z^{-1}za^{-1}a.$$

Then by the above theorem the one-relator inverse monoid

$$\text{Inv}\langle a, z, t \mid eazaz^{-1}a^{-1}za^{-1}z^{-1} = 1 \rangle$$

has undecidable word problem. This completes the proof. □

## Related work

### Other negative results

**Adjan (1966):** Proved the group of units of  $\text{Mon}\langle A \mid w = 1 \rangle$  is a one-relator group.

**Makanin (1966):** Proved that the monoid  $\text{Mon}\langle A \mid w_1 = 1, \dots, w_k = 1 \rangle$  has a finitely presented group of units (with  $k$  defining relations), and that  $M$  has decidable word problem if and only if its group of units also does.

In recent joint work with **Nik Ruškuc** we have shown:

- ▶ There is a one-relator inverse monoid  $\text{Inv}\langle A \mid w = 1 \rangle$  whose group of units is *not a one-relator group*.
- ▶ There is a finitely presented inverse monoid  $\text{Inv}\langle A \mid w_1 = 1, \dots, w_k = 1 \rangle$  whose group of units is *not finitely presented*.

### Some positive results

In recent joint work with **Igor Dolinka** we have shown the word problem is decidable for some new classes of  $\text{Inv}\langle A \mid w = 1 \rangle$  where  $w$  is a cyclically reduced and the maximal group image  $\text{Gp}\langle A \mid w = 1 \rangle$  is “low down” in the Magnus–Moldovanskii hierarchy.

# Open problems

## Problem

For which words  $w \in (A \cup A^{-1})^*$  does  $\text{Inv}\langle A \mid w = 1 \rangle$  have decidable word problem? In particular is the word problem always decidable when  $w$  is (a) **reduced** or (b) **cyclically reduced**?

# Open problems

## Problem

For which words  $w \in (A \cup A^{-1})^*$  does  $\text{Inv}\langle A \mid w = 1 \rangle$  have decidable word problem? In particular is the word problem always decidable when  $w$  is (a) **reduced** or (b) **cyclically reduced**?

## Problem

Characterise the one-relator groups with decidable submonoid membership problem.

## Problem

Characterise the one-relator groups with decidable rational subset membership problem.

## Problem

Is the subgroup membership problem decidable for one-relator groups?